

Orantes-Navarro CM, Herrera-Valdés R, Almaguer-López M, Brizuela-Díaz EG, Alvarado-Ascencio NP, Fuentes-de Morales EJ, et al. Chronic Kidney Disease in Children and Adolescents in Salvadoran Farming Communities: NefroSalva Pediatric Study (2009–2011). *MEDICC Rev.* 2016 Jan–Apr;18(1–2):15–21.

Page 19, Table 4. The second and third rows of data were misaligned by column. The corrected table is below.

Table 4: Prevalence of chronic kidney disease by stage, age group, sex and region

Variable (n)			CKD status				Total with CKD
			No CKD	Stage 1	Stage 2	Stages 3–5 n (stage)	
Total (1623)		n	1560	56	5	1 (3a) + 1 (5)	63
		%	96.1	3.5	0.3	0.1	3.9
		95% CI	95.2–97.1	2.6–4.3	0.0–0.6	0.0–0.3	2.9–4.8
Age	2–5 years (312)	n	296	14	2	0	16
		%	94.9	4.5	0.6		5.1
		95% CI	92.4–97.3	2.2–6.8	0.0–1.5		2.7–7.6
	6–12 years (756)	n	728	26	1	1(5)	28
		%	96.3	3.4	0.1	0.1	3.7
		95% CI	94.9–97.6	2.1–4.7	0.0–0.4	0.0–0.4	2.4–5.1
	13–17 years (555)	n	536	16	2	1 (3a)	19
		%	96.6	2.9	0.4	0.2	3.4
		95% CI	95.1–98.1	1.5–4.3	0.0–0.9	0.0–0.5	1.9–4.9
Sex	Male (800)	n	771	23	5	1 (3a)	29
		%	96.4	2.9	0.6	0.1	3.6
		95% CI	95.1–97.7	1.7–4.0	0.1–1.2	0.0–0.4	2.3–4.9
	Female (823)	n	789	33	0	1 (5)	34
		%	95.9	4.0		0.1	4.1
		95% CI	94.5–97.2	2.7–5.4		0.0–0.4	2.8–5.5
Region	Bajo Lempa (866)	n	831	31	4	0	35
		%	96.0	3.6	0.5		4.0
		95% CI	94.6–97.3	2.3–4.8	0.0–0.9		2.7–5.4
	Guayapa Abajo (412)	n	399	10	1	1 (3a) + 1 (5)	13
		%	96.8	2.4	0.2	0.5	3.2
		95% CI	95.2–98.5	0.9–3.9	0.0–0.7	0.0–1.2	1.5–4.8
	Las Brisas (345)	n	330	15	0	0	15
		%	95.7	4.3			4.3
		95% CI	93.5–97.8	2.2–6.5			2.2–6.5